


3.75 cm.





2.5 cm.

LIST OF TABLES

#

#

!	#	
Table	e	Page
3.1	Common types of dementia and their typical characteristics	17
4.1	Types of reactive oxygen species	29
4.2	Compared differences between necrosis and apoptosis	39
5.13	Cytotoxicity of H_2O_2 at 24 hours and the protection from mangosteen	65
	extract 200 and 400 μg/ml by Trypan blue assay	
5.14	Cytotoxicity of H ₂ O ₂ at various exposure times by MTT assay	67
5.15	Cytotoxicity from various concentrations of H ₂ O ₂ at 24 hours	69
	incubation measured by MTT assay	
5.16	Cytotoxicity from mangosteen extract on SK-N-SH cells various	70
	times by MTT assay	

3.75 cm.

2.5 cm.



LIST OF TABLES (cont.)

#

#

i	π		
Table		Page	
5.17	Cytotoxicity screening of mangosteen extract at various concentrations on	72	
	SK-N-SH cells at 24 hours incubation by MTT assay		
5.18	Protective effects from various incubation times of mangosteen	74	
	extract against H_2O_2 150, 300 μM induced SD-N-SH cells toxicity		
	measured by MTT assay		
5.19	Protective effects from mangosteen extract 25-800 µg/ml against	77	
	H_2O_2 cytotoxicity at 75, 150 and 300 μM on SK-N-SH cells		2.5 cm.
5.20	The percentages of cell population measured by flow cytometry	80	
6.1	Presumed caspase-3 activities from the protection of mangosteen	84	
	extract 200 $\mu g/ml~$ against $H_2O_2~150~\mu M~$ at 3,6 and 24 hour ixposure		
6.2	ROS production decreased from the protection of mangosteen extract	87	
	$200~\mu g/ml~$ against $150\text{-}600~$ $\mu M~$ H_2O_2 induced cell death during $0\text{-}180~$		
	min exposure		
6.3	ROS production after pretreated with mangosteen extract 200 µg/ml,	89	
	vitamin C 25-100 μM and vitamin E 5-25 μM against 150 μM		
	H_2O_2 exposusre		

3.75 cm.